

## PRELIMINARY COMMUNICATION

### EVALUATION OF INTERACTION BETWEEN VALPROATE AND BACLOFEN IN THE FORMALIN TEST IN MICE

*Mirosław Czuczwar<sup>1</sup>, Jacek Kiś<sup>1</sup>, Mariusz Świąder<sup>2</sup>, Waldemar A. Turski<sup>3</sup>,  
Krzysztof Przesmycki<sup>1,#</sup>*

<sup>1</sup>Second University Department of Anaesthesiology and Intensive Therapy, Medical University, Staszica 16, PL 20-081 Lublin, Poland, <sup>2</sup>Department of Pharmacology and Toxicology, Medical University, Jaczewskiego 8, PL 20-090 Lublin, Poland, <sup>3</sup>Department of Toxicology, Institute of Agricultural Medicine, Jaczewskiego 2, PL 20-090 Lublin, Poland

*Evaluation of interaction between valproate and baclofen in the formalin test in mice.*  
M. CZUCZWAR, J. KIŚ, M. ŚWIĄDER, W.A. TURSKI, K. PRZESMYCKI. Pol. J. Pharmacol., 2001, 53, 51–54.

Valproate and baclofen dose-dependently inhibited both phases of the formalin test. Combination of valproate and baclofen exerted the additive antinociceptive effect on both phases of the formalin test.

**Key words:** *baclofen, valproate, antinociception, formalin test, interaction, mice*